



PRONAMIDE 50WSP

PROPYZAMIDE | HERBICIDE

RESTRICTED USE PESTICIDE

Because Pronamide has produced tumors in laboratory animals, this product is for retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

GROUP

3

HERBICIDE

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR WILLOWOOD PRONAMIDE 50WSP BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Willowood Pronamide 50WSP and this supplement.

Willowood Pronamide 50WSP

EPA Reg. No. 87290-3

For Grass and Weed Control in Head Lettuce – Aerial Application
(For Use in Arizona and California)

KEEP OUT OF REACH OF CHILDREN
CAUTION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Willowood Pronamide 50WSP may be applied by aircraft for preemergence control of susceptible grasses and broadleaf weeds in head lettuce. Willowood Pronamide 50WSP must be applied at the dosage rate of 2.0 to 4.0 pounds of product (1.0 to 2.0 pounds active ingredient) per treated acre depending on soil type (refer to comments under Dosage below). Mix the specified amount of Willowood Pronamide 50WSP in 10 to 20 gallons of water per acre for aerial application. For aerial applications of Willowood Pronamide 50WSP on head lettuce, consult the section below for Rotation Crop Planting Information for plantback information. Avoid drift to all other crops and non-target areas.

Dosage

Willowood Pronamide 50WSP may be applied at the rate of 2 to 4 lbs. of product (1 to 2 lbs. active ingredient) per acre broadcast application. The dosage rate required is dependent on soil texture and method of irrigation. At rates specified on this label, Willowood Pronamide 50WSP may not be applied for weed control on highly organic (peat and muck) soils.

Follow the dosage instructions listed in chart below:

Lbs. of Willowood Pronamide 50WSP Per Broadcast Acre ¹			
Weeds	Dependable Rainfall or Overhead Irrigation	Less Dependable Rainfall or Furrow Irrigation	Soil Texture Group ²
susceptible annual grasses	2 - 3 (surface application)	3 - 4 (soil incorporation)	Coarse and medium Textured soils
broadleaf weeds	3 - 4 (surface application)	4 (soil incorporation)	Fine textured soils

¹ Reduce dosage rate accordingly for banded applications.

² Soil Texture Group

Coarse: sand, loamy sand, sandy loam

Medium: loam, silt loam, silt, sandy clay loam

Fine: silty clay loam, clay loam, sandy clay, silty clay, clay

Rotation Crop Planting Information

Follow the directions given below when rotation crops will be planted to areas previously treated with Willowood Pronamide 50WSP:

Waiting Period in Days Before Planting the Crops Indicated (1):

Amount of Willowood Pronamide 50WSP Applied per Planted Acre	Root and Tuber Vegetables	Leafy Vegetables (except Brassica Vegetables), Crop Group 4 (2)	Barley, Buckwheat, Field Corn, Millet, Oats, Pop-corn, Rice, Rye, Sorghum (milo), Teosinte, Triticale, Wheat, and Wild Rice
1.0 lb	90	30	180
2.0 lb	90	30	180
3.0 lb	90	30	180
4.0 lb	90	30	180

(1) There are no plant back restrictions for Willowood Pronamide 50WSP when rotating to artichokes, grapes, berry fruits, pome fruits or stone fruits.

(2) Crop Group 4 as defined under 40CFR 180.41.

Whether Willowood Pronamide 50WSP is bed-topped, banded, or broadcast, the beds must be knocked down and the field cross-disced before rotation crops other than artichokes, head lettuce, endive, radicchio or escarole are planted.

Where the Willowood Pronamide 50WSP treatment is to be followed by a rotation crop within 180 days of application, bed-topped or banded applications are suggested.

Read the “**CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**” in the label booklet for Willowood Pronamide 50WSP before using.



WILLOWOOD USA

Willowood, LLC
1600 NW Garden Valley Blvd. #120
Roseburg, OR 97471

EPA 20110218