



# PRONAMIDE 50WSP

PROPYZAMIDE | HERBICIDE

## RESTRICTED USE PESTICIDE

Because Pronamide has produced tumors in laboratory animals, this product is for retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

GROUP 3 HERBICIDE

### SUPPLEMENTAL LABELING

## Willowood Pronamide 50 WSP

EPA Reg. No. 87290-3

24(c) Special Local Need Registration SLN OR-110002

**Weed Control in Fall, Carbon-Seeded Grasses (Orchardgrass, Tall Fescue, Perennial Ryegrass) Grown for Seed**  
(For Distribution and Use Only in the State of Oregon)

This Special Local Need label must only be used with Willowood Pronamide 50 WSP (EPA Reg. No. 87290-3), and can not be used with Willowood Pronamide 50 (EPA Reg. No. 82542-22-87290).

**Expiration Date:** This label is valid until December 31, 2016, or until otherwise amended, withdrawn, canceled, or suspended.

**ATTENTION**  
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Willowood Proamide 50 WSP and this Section 24c label before applying. Do not use in manner inconsistent with applicable directions, restrictions, and precautions provided on the label affixed to the container for Willowood Proamide 50 WSP and this Section 24c label. Use of Willowood Proamide 50 WSP according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Willowood Proamide 50 WSP.

**ACTIVE INGREDIENT:**

Pronamide: 3,5-dichloro-N-(1,1-dimethyl-2-propynyl) benzamide .....50.0%

**OTHER INGREDIENTS:** .....50.0%

**TOTAL:** .....100.0%

# KEEP OUT OF REACH OF CHILDREN CAUTION

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye irritation. Avoid contact with eyes or clothing.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### General Information

Willowood Pronamide 50 WSP has been demonstrated to be effective for the control and/or suppression of undesirable seedling grasses.

**NOTE:** *Due to the close proximity of native prairie remnants to agricultural areas and the potential for these areas to be adversely affected by herbicides through drift or possible runoff/soil movement, Willowood Pronamide 50 WSP should not be applied directly to native prairie habitats. It is recommended that dosage rates be maintained at the lower end of the use rate range (if effective seedling control can be achieved) in a further effort to lessen potential impacts to endangered species.*

#### Weeds Controlled (Suppression)

Willowood Pronamide 50 WSP will provide preemergence and postemergence control/suppression of the following seedling weeds when used according to label directions:

Bromus Species	<i>Bromus species</i>
Wild Oat	<i>Avena species</i>
Annual Ryegrass	<i>Lolium multiflorum</i>
Volunteer Orchardgrass	<i>Dactylis glomerata</i>

#### Application Rate

Prior to application of Willowood Pronamide 50 WSP, ensure that an adequate activated carbon band (one inch wide or greater) to absorb the herbicide applied is present on a well-prepared, firmly packed seedbed that is free of clods. After seeding/carbon banding operation, apply a single application of Willowood Pronamide 50 WSP at 0.25 to 0.5 lb (0.125 to 0.25 lb ai) per acre tank mixed with a reduced rate of diuron. Application must be made before the emergence of the seeded crop and before rainfall dissipates the carbon band applied. Consult your local Extension Service agent or crop consultant for specific application rate recommendations for Willowood Pronamide 50 WSP, diuron, and the activated charcoal on a field by field basis. **The grower assumes all risks of crop injury or stand loss when unforeseen environmental conditions (such as heavy rain soon after application), poor seedbed preparation or application rates/positioning, result in crop damage.** The greatest risk of injury or stand loss occurs when applications are made to poorly drained soils and heavy rains occur soon after application.

Do not apply Willowood Pronamide 50 WSP to sandy or gravelly areas or fields under stress due to drought, flooding, excessive fertilizer, soil salts, cold temperatures, hail, injury from previously applied pesticides or injury due to insects or diseases, as excessive injury may result.

#### Specific Use Restrictions

The following use restrictions must be observed when Willowood Pronamide 50 WSP is used as recommended on this label:

- **Chemigation:** Do not apply this product through any type of irrigation system.
- Read and observe all label directions before using. All applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered product label must be followed.
- **Do not** graze livestock in treated fields within 180 days of application.

- **Do not** apply Willowood Pronamide 50 WSP within 180 days of harvesting grass hay.
- Willowood Pronamide 50 WSP should only be applied using ground application equipment.



Willowood, LLC  
1600 NW Garden Valley Blvd. #130  
Roseburg, OR 97471

Specimen