



# PRONAMIDE 50WSP

PROPYZAMIDE | HERBICIDE

## RESTRICTED USE PESTICIDE

Because Pronamide has produced tumors in laboratory animals, this product is for retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

GROUP 3 HERBICIDE

### SUPPLEMENTAL LABELING

## Willowood Pronamide 50 WSP

EPA Reg. No. 87290-3

24(c) Special Local Need Registration SLN OR-110001

### Weed Control in Established Orchardgrass Grown for Seed (For Distribution and Use Only in the State of Oregon)

This Special Local Need label must only be used with Willowood Pronamide 50 WSP (EPA Reg. No. 87290-3), and can not be used with Willowood Pronamide 50 (EPA Reg. No. 82542-22-87290).

**Expiration Date:** This label is valid until December 31, 2016, or until otherwise amended, withdrawn, canceled, or suspended.

**ATTENTION**  
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Willowood Proamide 50 WSP and this Section 24c label before applying. Do not use in manner inconsistent with applicable directions, restrictions, and precautions provided on the label affixed to the container for Willowood Proamide 50 WSP and this Section 24c label. Use of Willowood Proamide 50 WSP according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Willowood Proamide 50 WSP.

**ACTIVE INGREDIENT:**

Pronamide: 3,5-dichloro-N-(1,1-dimethyl-2-propynyl) benzamide.....50.0%

**OTHER INGREDIENTS:** .....50.0%

**TOTAL:** .....100.0%

# KEEP OUT OF REACH OF CHILDREN CAUTION

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye irritation. Avoid contact with eyes or clothing.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### General Information

Willowood Pronamide 50 WSP is effective for the control and/or suppression of undesirable seedling grasses.

**NOTE:** Due to the close proximity of native prairie remnants to agricultural areas and the potential for these areas to be adversely affected by herbicides through drift or possible runoff/soil movement, Willowood Pronamide 50 WSP should not be applied directly to native prairie habitats. It is recommended that dosage rates be maintained at the lower end of the use rate range (if effective seedling control can be achieved) in a further effort to lessen potential impacts to endangered species.

#### Weeds Controlled (Suppression)

Willowood Pronamide 50 WSP will provide preemergence and postemergence control/suppression of the following seedling weeds when used according to label directions:

Bromus Species	<i>Bromus species</i>
Wild Oat	<i>Avena species</i>
Annual Ryegrass	<i>Lolium multiflorum</i>
Volunteer Orchardgrass	<i>Dactylis glomerata</i>

#### Application Rate

Apply a single application of Willowood Pronamide 50 WSP at a rate of 0.5 to 0.75 lb (0.25 to 0.375 lb ai) per acre for preemergence and postemergence control/suppression of undesirable seedling grasses in established orchardgrass fields that have produced at least one seed crop. Application should be made during the fall or early winter months (November through mid-January), once the fall rains have moistened the soil and the soil temperature is 55°F or less. Applications must be completed by mid-January. Do not apply more than 0.75 lb (0.375 lb ai) of Willowood Pronamide 50 WSP per acre during any one season.

Do not apply Willowood Pronamide 50 WSP to sandy or gravelly areas or fields under stress due to drought, flooding, excessive fertilizer, soil salts, cold temperatures, hail, injury from previously applied pesticides or injury due to insects or diseases, as excessive injury may result.

#### Specific Use Restrictions

The following use restrictions must be observed when Willowood Pronamide 50 WSP is used as recommended on this label:

- **Chemigation:** Do not apply this product through any type of irrigation system.
- Read and observe all label directions before using. All applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered product label must be followed.
- **Do not** graze livestock in treated fields within 180 days of application.
- **Do not** apply Willowood Pronamide 50 WSP within 180 days of harvesting grass hay.
- Willowood Pronamide 50 WSP should only be applied using ground application equipment.



**WILLOWOOD USA**

Willowood, LLC  
1600 NW Garden Valley Blvd. #130  
Roseburg, OR 97471

Specimen