

Contains the  
same active as

**ELITE® DF**



**WILLOWOOD USA** 

# TEBUCON 45DF

TEBUCONAZOLE | FUNGICIDE

## BENEFITS

- Cost effective disease treatment
- Dry flowable (DF) formulation is easy to handle
- Mode of action (DMI) works well in a resistance management program
- Absorbed through leaves and blossoms
- Steady, reliable disease control
- Curative and preventative disease activity
- Excellent crop safety

**Tebucon 45DF** fungicide provides highly effective, economical control of powdery mildew, black rot, leafspot, and brown rot in stone fruits and grapes.

A locally systemic fungicide, Tebucon 45DF fungicide forms the basis of resistance management programs.

Tebucon 45DF fungicide also offers flexible use rates and a short 12-hour reentry interval, plus a 0-day pre-harvest interval on stone fruit and a 14-day pre-harvest interval on grapes.

Tebuconazole is used on the following crops—cherry, peach, nectarine and grapes.



## GRAPES:

- Tebucon 45DF fungicide provides unsurpassed protection against powdery mildew, which can cause adverse effects on grape quality.
- Tebucon 45DF fungicide enhances control of Botrytis bunch rot, which can lead to entire bunches of infected, blemished grapes
- Tebucon 45DF fungicide is a sterol demethylation inhibitor (DMI), which is a foundation for resistance management when used in combination with a strobilurin fungicide.

## STONE FRUITS :

- Tebucon 45DF fungicide effectively controls all forms of brown rot in stone fruit from first bloom to pre-harvest
- Tebucon 45DF fungicide enters stone fruit blossoms and leaves and is absorbed into plant tissue, resulting in more consistent disease control
- Tebucon 45DF fungicide also effectively controls blossom blight and fruit rot on stone fruit varieties

## TEBUCON 45DF FUNGICIDE

### REGISTERED USES :

This crop list is a compilation from all Federal, State and local labels for this product including limited use Section 18 emergency use, 24(c) and 2EE labels. Please consult the label to ensure your intended use is registered in your area.

- ✓ Banana
- ✓ Nectarine
- ✓ Cherry
- ✓ Peach
- ✓ Grape
- ✓ Plantain

### WHEN TO USE:

Tebucon 45DF fungicide may be applied up to and including the day of harvest (0-day PHI) on selected crops. Do not apply more than 3 lb of product per acre per crop season.

### HOW TO USE:

Always read and follow all label directions.

### ACTIVE INGREDIENTS:

Tebuconazole	45%
Other Ingredients	55%

EPA Reg. No. 87290-12

EPA Est. No. 39578-TX-001



**WILLOWOOD USA** 

### CORPORATE OFFICE:

1600 NW Garden Valley Blvd #120 • Roseburg, OR 97471  
541-679-9963 • 877-679-9963 • Fax: 541-679-4650  
[www.WillowoodUSA.com](http://www.WillowoodUSA.com)