

# RESTRICTED USE PESTICIDE

Because Pronamide has produced tumors in laboratory animals, this product is for retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

# WILLOWOOD PRONAMIDE 3.3SC

## SUPPLEMENTAL LABELING

### Willowood Pronamide 3.3SC

EPA Reg. No. 87290-22

24(c) Special Local Need Registration SLN OR-xxxxxx

### Weed Control in Fall, Carbon-Seeded Grasses (Orchardgrass, Tall Fescue, Perennial Ryegrass) Grown for Seed (For Distribution and Use Only in the State of Oregon)

**Expiration Date:** This label is valid until December 31, 2017, or until otherwise amended, withdrawn, canceled, or suspended.

#### ATTENTION

- This label and the federal label for this product must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, and precautions on this supplemental label and the main EPA-registered label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### ACTIVE INGREDIENT:

Pronamide: 3,5-dichloro-N-(1,1-dimethyl-2-propynyl) benzamide..... 35.6%

**OTHER INGREDIENTS:** ..... 64.4%

**TOTAL:**..... 100.0%

Contains 3.3 lbs. of active ingredient per gallon.

## KEEP OUT OF REACH OF CHILDREN CAUTION

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye irritation. Avoid contact with eyes or clothing.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### Product Information

Willowood Pronamide 3.3SC has been demonstrated to be effective for the control and/or suppression of undesirable seedling grasses.

**NOTE:** Due to the close proximity of native prairie remnants to agricultural areas and the potential for these areas to be adversely affected by herbicides through drift or possible runoff/soil movement, Willowood Pronamide 3.3SC should not be applied directly to native prairie habitats. It is recommended that dosage rates be maintained at the lower end of the use rate range (if effective seedling control can be achieved) in a further effort to lessen potential impacts to endangered species.

### Weeds Controlled (Suppression)

Willowood Pronamide 3.3SC will provide preemergence and postemergence control/suppression of the following seedling weeds when used according to label directions:

|                        |                           |
|------------------------|---------------------------|
| Bromus Species         | <i>Bromus species</i>     |
| Wild Oat               | <i>Avena species</i>      |
| Annual Ryegrass        | <i>Lolium multiflorum</i> |
| Volunteer Orchardgrass | <i>Dactylis glomerata</i> |

### Application Rate

Prior to application of Willowood Pronamide 3.3SC, ensure that an adequate activated carbon band (one inch wide or greater) to absorb the herbicide applied is present on a well-prepared, firmly packed seedbed that is free of clods. After seeding/carbon banding operation, apply a single application of Willowood Pronamide 3.3SC at 5.0 to 10.0 fl. oz. (0.125 to 0.25 lb ai) per acre tank mixed with a reduced rate of diuron. Application must be made before the emergence of the seeded crop and before rainfall dissipates the carbon band applied. Consult your local Extension Service agent or crop consultant for specific application rate recommendations for Willowood Pronamide 3.3SC, diuron, and the activated charcoal on a field by field basis. **The grower assumes all risks of crop injury or stand loss when unforeseen environmental conditions (such as heavy rain soon after application), poor seedbed preparation or application rates/positioning, result in crop damage.** The greatest risk of injury or stand loss occurs when applications are made to poorly drained soils and heavy rains occur soon after application.

Do not apply Willowood Pronamide 3.3SC to sandy or gravelly areas or fields under stress due to drought, flooding, excessive fertilizer, soil salts, cold temperatures, hail, injury from previously applied pesticides or injury due to insects or diseases, as excessive injury may result.

### Specific Use Restrictions

The following use restrictions must be observed when Willowood Pronamide 3.3SC is used as recommended on this label:

- **Chemigation:** Do not apply this product through any type of irrigation system.
- Read and observe all label directions before using. All applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered product label must be followed.
- **Do not** graze livestock in treated fields within 180 days of application.
- **Do not** apply Willowood Pronamide 3.3SC within 180 days of harvesting grass hay.
- Willowood Pronamide 3.3SC should only be applied using ground application equipment.



Willowood, LLC  
1600 NW Garden Valley Blvd. #120  
Roseburg, OR 97471