

RESTRICTED USE PESTICIDE

Because Pronamide has produced tumors in laboratory animals, this product is for retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

WILLOWOOD PRONAMIDE 3.3SC

SUPPLEMENTAL LABELING

Willowood Pronamide 3.3SC

EPA Reg. No. 87290-22

24(c) Special Local Need Registration SLN OR-xxxxxx

Weed Control in Established Orchardgrass Grown for Seed (For Distribution and Use Only in the State of Oregon)

Expiration Date: This label is valid until December 31, 2017, or until otherwise amended, withdrawn, canceled, or suspended.

ATTENTION

- This label and the federal label for this product must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, and precautions on this supplemental label and the main EPA-registered label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ACTIVE INGREDIENT:

Pronamide: 3,5-dichloro-N-(1,1-dimethyl-2-propynyl) benzamide..... 35.6%

OTHER INGREDIENTS: 64.4%

TOTAL:..... 100.0%

Contains 3.3 lbs. of active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye irritation. Avoid contact with eyes or clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Product Information

Willowood Pronamide 3.3SC is effective for the control and/or suppression of undesirable seedling grasses.

NOTE: Due to the close proximity of native prairie remnants to agricultural areas and the potential for these areas to be adversely affected by herbicides through drift or possible runoff/soil movement, Willowood Pronamide 3.3SC should not be applied directly to native prairie habitats. It is recommended that dosage rates be maintained at the lower end of the use rate range (if effective seedling control can be achieved) in a further effort to lessen potential impacts to endangered species.

Weeds Controlled (Suppression)

Willowood Pronamide 3.3SC will provide preemergence and postemergence control/suppression of the following seedling weeds when used according to label directions:

Bromus Species	<i>Bromus species</i>
Wild Oat	<i>Avena species</i>
Annual Ryegrass	<i>Lolium multiflorum</i>
Volunteer Orchardgrass	<i>Dactylis glomerata</i>

Application Rate

Apply a single application of Willowood Pronamide 3.3SC at a rate of 10.0 to 15.0 fl. oz. (0.25 to 0.375 lb ai) per acre for preemergence and postemergence control/suppression of undesirable seedling grasses in established orchardgrass fields that have produced at least one seed crop. Application should be made during the fall or early winter months (November through mid-January), once the fall rains have moistened the soil and the soil temperature is 55°F or less. Applications must be completed by mid-January. Do not apply more than 15 fl. oz. (0.375 lb ai) of Willowood Pronamide 3.3SC per acre during any one season.

Do not apply Willowood Pronamide 3.3SC to sandy or gravelly areas or fields under stress due to drought, flooding, excessive fertilizer, soil salts, cold temperatures, hail, injury from previously applied pesticides or injury due to insects or diseases, as excessive injury may result.

Specific Use Restrictions

The following use restrictions must be observed when Willowood Pronamide 3.3SC is used as recommended on this label:

- **Chemigation:** Do not apply this product through any type of irrigation system.
- Read and observe all label directions before using. All applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered product label must be followed.
- **Do not** graze livestock in treated fields within 180 days of application.
- **Do not** apply Willowood Pronamide 3.3SC within 180 days of harvesting grass hay.
- Willowood Pronamide 3.3SC should only be applied using ground application equipment.



WILLOWOOD USA

Willowood, LLC
1600 NW Garden Valley Blvd. #120
Roseburg, OR 97471